

University of Groningen

Nieuwe typen van onverzadigde ethers, thioethers en selenoethers

Brandsma, Lambert

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1961

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Brandsma, L. (1961). *Nieuwe typen van onverzadigde ethers, thioethers en selenoethers*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.


SUMMARY

In this thesis the preparation of some new types of unsaturated ethers, thioethers and selenoethers is described.

Chapter I deals with the preparation of di(alk-1-enyl)ethers, -thioethers and -selenoethers $RCH=CH-X-CH=CHR$ (X is O, S or Se, R is H or alkyl). The oxygen-ethers were obtainable in a simple manner, starting from aliphatic aldehydes. The latter compounds were first converted into 1,1-dichlorodialkyl ethers (I) by means of anhydrous hydrogen chloride; the dichloroethers in turn were heated with diethylaniline, which produced the di(alk-1-enyl)ethers (II):





The analogous thioethers and selenoethers (IIa) were prepared by heating 1,1'-dichlorodialkylthioethers or -selenoethers (Ia) with diethylaniline. The latter compounds were accessible by passing a mixture of gaseous HCl and H_2S or H_2Se in the mol. proportion 2 : 1 into aldehydes:



The overall yields were reasonable.


In chapter II the synthesis of compounds, containing the system $-C\equiv C-X-CH\equiv C-$ (X is O, S or Se) is described. The synthetic routes are apparent from the reaction schemes A-C:



In the first scheme sulfur dichloride is added to an 1-alkyne; the adduct furnishes the di(alk-1-ynyl)thioethers, on treating with two equivalents of LiNH_2 in liquid ammonia.

In the second scheme 2, 2'-dibromodivinylthioether (obtained from divinylthioether) is converted into the disodio-compound of diethynylthioether, which can be alkylated to the dialkynylthioethers (IV) or hydrolyzed to diethynylthioether (III).

The formation of the extremely unstable dialkynylethers $(\text{RC}\equiv\text{C})_2\text{O}$ of which only dipropynylether $(\text{CH}_3\text{C}\equiv\text{C})_2\text{O}$ has been isolated, proceeded as follows:



The first representative $(\text{HC}\equiv\text{C})_2\text{O}$ seemed to be unstable even at -70° .